
Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.



62.81

CLINT McDADE & SONS



Rivermont Orchids

June 1, 1948

SPECIAL OFFERING

MATURE PLANTS OF CATTLEYA SPECIES

These plants are well established in 5" to 8" pots. The price is determined by the size and weight of the plant, together with the number of new leads that are growing. Minimum standards are as follows:

Small ---- 5 to 7 Bulbs and Leaves
Medium ---- 6 to 8 " " "
Large ---- 7 to 10 " " "

Prices are F.O.B. Chattenooga. Plants will be shipped by Railway Express unless otherwise directed.

	PLANT SIZE	PRICE
CATTLEYA TRIANAE	Small	\$5.00
	Medium	7.50
	Large	10.00

Normal blooming season is December through February. The flowers have light lavender petals and sepals and a dark lavender lip. One of the better known species.

CATTLEYA SCHRODERAE	Small	5.00
	Medium	7.50
	Large	10.00

Normal blooming season is February through April. The flowers are of varying shades of pink with a bright yellow throat. Although the blooms are not as large as those of C. Trianae, this genera is floriferous and will usually average three or more flowers per spike.

200,88 MAY

CLINT McDADE & SONS

Rivermont Onhide

STEPHNOT MIATRUOM JAMENS (binvelope some SICH & SIEDA amelgale?

June 1, 1948

SERVINE CONTROLL

MATCHE PLANTS OF CATCLEY SPECIES

These plants are well setablished in 5" to 8" pots. The price is detailed to the the state of the plant, to the the number of new leads the the number of new leads to the crowing. Whitman standards are so vollows:

Small ----- 5 to V Bulbu and Leaves
Medium ---- 6 to 8 " " "
Large ---- 7 to 10 " " "

Prince or F.O.B. Chattenooga. Flemts will be shipped by Railway Express unless otherwise directed.

TIMES SAMATOR AVELTAGE CO.O. DATES

Morriel blooming season to Locember through February. The flowers have light lavender potals and sepals and a durk lavender lig. One of the better such appeals

CATTLEYA SCHIODERAL

Amall E.00 Wedium 7.50 Large 10.00

Normal blooming member is Pebruary through April. The flowers are of verylag shedes of pink with a bright yellow throat. Although the blooms are not as large as those of C. Trianne, this genera is floriferous and will usually everage three or more flowers per apike.